

SuperPlex[™] Premix

Catalog Nos.	Amount	Lot
638543	200 rxns	Spec
638544	1,000 rxns	Spec

Lot Number Specified on product label. Specified on product label.

Description

SuperPlex Premix is a 2X master mix containing a 5' to 3' exonuclease-deficient *Taq* polymerase and the TaqStart® Antibody, a monoclonal antibody that inhibits the activity of *Taq* and *Taq*-derived polymerases at ambient temperatures during PCR reaction assembly. The presence of the TaqStart Antibody in the polymerase mix allows automatic hot-start PCR. Enough premix is supplied for 200 or 1,000 reactions of 50 μ l each.

Package Contents

 638543
 638544

 (200 rxns)
 (1,000 rxns)

 5 x 1 ml
 5 x 5 ml
 SuperPlex Premix

Storage Conditions

• Store at -20°C.

Expiration Date

• Specified on product label.

Shipping Conditions

• Dry ice

Product Documents

Documents for our products are available for download at <u>takarabio.com/manuals</u> The following documents apply to this product:

• SuperPlex Premix Protocol-At-A-Glance

Certificate of Analysis

SuperPlex Premix

Quality Control Data

Raw-Material Quality Control

SuperPlex Premix was tested for enzymatic activity and PCR performance. Endonuclease, exonuclease, and DNA contamination assays were also performed.

Functional Quality Control

Amplification from a cDNA template:

SuperPlex Premix was tested in a 50 μ l PCR reaction using 5 μ l of Marathon®-Ready Human Placenta cDNA (Cat. No. 639311) as template and primers (0.2 μ M each) specific for a 1.3 kb fragment from the transferrin receptor gene. Cycling conditions were set at:

95°C 1 min 30 cycles: 95°C 15 sec 68°C 1.5 min 68°C 3 min

5 μ l of the PCR product was run on a 1.2% TAE/agarose gel to confirm the presence of a 1.3 kb product with minimal background. PCR product concentration was measured by fluorometry. The yield was determined to be >5 ng/ μ l.

Amplification from a genomic DNA template

SuperPlex Premix was tested in a 50 μ l PCR reaction using 100 ng of calf thymus genomic DNA as template and primers (0.4 μ M each) specific for a 407-bp fragment of the bovine pancreatic trypsin inhibitor (BPTI) gene. Cycling conditions were set at:

94°C		3 min	
30 cycle	s:		
	94°C	30 sec	
	68°C	1.5 min	
68°	С	3 min	<u> </u>

5 μ l of PCR product was run on a 1.2% TAE/agarose gel to confirm the presence of a 407 bp product with minimal background. PCR product concentration was measured by fluorometry. The yield was determined to be >15 ng/µl.

It is certified that this product meets the above specifications, as reviewed and approved by the Quality Department.



SuperPlexTM Premix

CATALOG NOS.

638543 & 638543

NOTICE TO PURCHASER:

Our products are to be used for **Research Use Only**. They may not be used for any other purpose, including, but not limited to, use in humans, therapeutic or diagnostic use, or commercial use of any kind. Our products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products or to provide a service to third parties without our prior written approval.

Your use of this product is also subject to compliance with the licensing requirements, listed below if applicable, and described on the product's web page at <u>http://www.takarabio.com</u>. It is your responsibility to review, understand and adhere to any restrictions imposed by these statements.

TRADEMARKS:

©2023 Takara Bio Inc. All Rights Reserved.

All trademarks are the property of Takara Bio Inc. or its affiliate(s) in the U.S. and/or other countries or their respective owners. Certain trademarks may not be registered in all jurisdictions.

Takara Bio USA, Inc.					
2560 Orchard Parkway, Sar	1 Jose, CA 95131, USA	4			
U.S. Technical Support: technical_support@takarabio.com					
United States/Canada	Asia Pacific	Europe	Japan	6/14/2023	
800.662.2566	+1.650.919.7300	+33.(0)1.3904.6880	+81.(0)77.565.6999		