

# **Cloning Enhancer**

Catalog Nos.	Amount	Lot Number
639613 (Not sold separately; sold as a part of 639633)	50 μl	Specified on product label.
639614 (Not sold separately; sold as a part of 639634)	100 μl	Specified on product label.
639615	200 μl	Specified on product label.
638926	500 μl	Specified on product label.

## **Description**

Cloning Enhancer removes background plasmid DNA and PCR residue, eliminating the need for PCR insert purification prior to cloning. Cloning Enhancer-treated PCR samples yield significantly more recombinant clones. Since treatment with Cloning Enhancer occurs in the same tube as the PCR reaction, it is less likely to result in UV damage or nicking. Avoiding a separate cleanup process also minimizes the risk of losing PCR products during purification.

## **Package Contents**

639613	<u>639614</u>	<u>639615</u>	<u>638926</u>	
(25  rxns)	(50  rxns)	(100  rxns)	(250  rxns)	
50 µl	2 x 50 ul	4 x 50 ul	500 ul	Cloning Enhancer

## **Storage Conditions**

Store at  $-20^{\circ}$ C.

## **Expiration Data**

Specified on product label.

### **Shipping Conditions**

Dry ice

### **Product Documents**

Documents for our products are available for download at takarabio.com/manuals The following documents apply to this product:

- Cloning Enhancer Protocol-At-A-Glance
- In-Fusion® Snap Assembly Master Mix User Manual

## Quality Control Data

An aliquot of this lot of Cloning Enhancer was tested using the components of the In-Fusion Snap Assembly Master Mix (Takara Bio, Cat. No. 638947). An aliquot of Cloning Enhancer was mixed with a 2.0 kb Unpurified Control Insert and incubated at 37°C for 15 min, followed by 80°C for 15 min. After Cloning Enhancer treatment, the Unpurified Control Insert was cloned into the linearized pUC19 Vector as described in the In-Fusion Snap Assembly Master Mix User Manual. The cloning reaction was used to transform chemically competent Stellar<sup>TM</sup> Competent Cells (Takara Bio, Cat. No. 636763) (> 5 x 10<sup>8</sup> cfu/µg). After 1 hr of growth, an aliquot of the transformed culture was plated onto an LB Amp 100/X-gal/IPTG plate. At least 2,500 colonies were observed for the cloning reaction.

It is certified that this product meets the above specifications, as reviewed and approved by the Quality Department.

2560 Orchard Parkway, San Jose, CA 95131, USA U.S. Technical Support: technical support@takarabio.com



8/10/2023

# **Cloning Enhancer**

## CATALOG NO.

638926, 639613, 639614 & 639615

#### NOTICE TO PURCHASER:

Our products are to be used for Research Use Only. They may not be used for any other purpose, including, but not limited to, use in humans, therapeutic or diagnostic use, or commercial use of any kind. Our products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products or to provide a service to third parties without our prior written approval.

Your use of this product is also subject to compliance with the licensing requirements, listed below if applicable, and described on the product's web page at http://www.takarabio.com. It is your responsibility to review, understand and adhere to any restrictions imposed by these statements.

### TRADEMARKS:

### ©2023 Takara Bio Inc. All Rights Reserved.

All trademarks are the property of Takara Bio Inc. or its affiliate(s) in the U.S. and/or other countries or their respective owners. Certain trademarks may not be registered in all jurisdictions.

Takara Bio USA, Inc.

2560 Orchard Parkway, San Jose, CA 95131, USA U.S. Technical Support: technical\_support@takarabio.com

800.662.2566 +1.650.919.7300 +81.(0)77.565.6999 +33.(0)1.3904.6880

United States/Canada Asia Pacific **Europe** Japan