



		XhoI									EcoRI
	End of AcGFP1		BglII							BstBI	
2902	CTG TAC AAG	TCC GGA CTC	AGA TCT CGA	GCT CAA GCT	TCG AAT TCT	GCA					
	SalI		ApaI	BamHI			Stop	Stop	Stop		
2950	GTC GAC GGT	ACC GCG GGC	CCG GGA TCC	ACC GGA TCT	AGA TAA	CTG ATC					

pRetroQ-AcGFP1-C1 Vector Map and Multiple Cloning Site (MCS).

Description

pRetroQ-AcGFP1-C1 is a high-titer, self-inactivating retroviral vector that facilitates efficient delivery and expression of AcGFP1, as well as C-terminal fusions of AcGFP1, to target cells. AcGFP1 is the green fluorescent protein from *Aequorea coerulea* (AcGFP1; Excitation maximum = 475 nm; emission maximum = 505 nm).

The AcGFP1 coding sequence contains silent base changes, which correspond to human codon-usage preferences (1). The multiple cloning site (MCS) in pRetroQAcGFP1-C1 follows the AcGFP1 coding sequence. Genes cloned into the MCS are expressed as fusions to the C-terminus of AcGFP1 when they are in the same reading frame as AcGFP1 and there are no intervening stop codons.

The RetroQ retroviral vector backbone incorporates several unique features. This vector contains a puromycin resistance cassette (Puro^r) driven by the PGK promoter (P_{PGK}) for selection of positively-infected cells (2). The hybrid 5' long terminal repeats (LTR) consists of the CMV type I enhancer and the murine sarcoma virus (MSV) promoter. This vector demonstrates high levels of transcription in HEK 293-based packaging cell lines due, in part, to the presence of adenoviral E1A (3-6) in these cells. The self-inactivating feature of the vector is provided by a deletion in the 3' LTR enhancer region (U3). During reverse transcription of the retroviral RNA, the inactivated 3' LTR is copied and replaces the 5' LTR CMV enhancer sequences. This can reduce the phenomenon known as promoter interference (7) and allow more efficient expression.

(PR983321; published 17 August 2009)



Clontech

United States/Canada
800.662.2566

Asia Pacific
+1.650.919.7300

Europe
+33.(0)1.3904.6880

Japan
+81.(0)77.543.6116

Clontech Laboratories, Inc.
A Takara Bio Company
1290 Terra Bella Ave.
Mountain View, CA 94043
Technical Support (US)
E-mail: tech@clontech.com
www.clontech.com

Additionally, the viral genomic transcript contains the necessary viral RNA processing elements, including the LTRs, packaging signal (Ψ^+), and tRNA primer-binding site. pRetroQ-AcGFP1-C1 contains a bacterial origin of replication, an *E. coli* Amp^r gene for propagation and selection in bacteria, and an SV40 origin for replication in mammalian cells expressing the SV40 large T antigen.

Use

pRetroQ-AcGFP1-C1 is designed to efficiently deliver and express fusions to the C terminus of AcGFP1 into primary cells or cells that are difficult to transfect. Fusions to the C terminus of AcGFP1 retain the fluorescent properties of the native protein, allowing the *in vivo* localization of the fusion protein. The target gene should be cloned into pRetroQ-AcGFP1-C1 so that it is in frame with the AcGFP1 coding sequences, with no intervening in-frame stop codons. The inserted gene should include the initiating ATG codon. The recombinant AcGFP1 vector can be infected or transfected into mammalian cells. If required, stable transformants can be selected using Puromycin. pRetroQ-AcGFP1-C1 can also be used simply to express AcGFP1 in a cell line of interest (e.g., as an infection marker).

Once pRetroQ-AcGFP1-C1 is transfected into a packaging cell line (such as the RetroPack™ PT67 Cell line (Cat. No. 631510), AmphiPack™-293 (Cat. No. 631505), EcoPack™ 2-293 (Cat. No. 631507), Pantropic Expression System (Cat. No. 631512), or Retro-X™ Universal Packaging System (Cat. No. 631530)), RNA from the vector is packaged into non-infectious, replication-incompetent retroviral particles, since pRetroQ-AcGFP1-C1 lacks the structural genes (*gag*, *pol*, and *env*) necessary for particle formation and replication. These genes, however, are stably integrated as part of the packaging cell genome. Once a high-titer supernatant is produced, these retroviral particles can infect target cells and transmit the gene of interest but cannot replicate within these cells due to the absence of viral structural genes. The separate introduction and integration of the structural genes into the packaging cell line minimizes the chances of producing replication-competent virus due to recombination events during cell proliferation.

Location of features

- 5' LTR (CMV/MSV): 1–727
 - Cytomegalovirus (CMV)/mouse sarcoma virus (MSV) hybrid promoter: 1–511
 - R region: 583–653
 - U5 region: 654–727
- Ψ^+ (extended packaging signal): 757–1566
- P_{CMVIE} (cytomegalovirus immediate early promoter): 1582–2170
- AcGFP1 (*Aequorea coerulescens* green fluorescent protein gene): 2194–2910
 - Start codon: (ATG): 2194–2196
 - Last codon: 2908–2910
- MCS (multiple cloning site): 2920–2976
- P_{PGK} (PGK promoter): 3000–3508
- Puro^r (Puromycin resistance gene): 3529–4128
- 3' LTR (MMLV; deletion in U3): 4313–4745
 - PolyA signal: 4571–4586
- P_{SV40} (SV40 promoter): 5025–5292
- SV40 origin of replication: 5246–5311
- ColE1 origin of replication: 5632
- Amp^r (ampicillin resistance gene; β -lactamase): 6392–7252 (complementary)

Propagation in *E. coli*

- Suitable host strains: DH5 α ™, Fusion Blue, and other general purpose strains.
- Selectable marker: plasmid confers resistance to ampicillin (100 μ g/ml) in *E. coli* hosts.
- *E. coli* replication origin: ColE1
- Copy number: low

NOTE: The viral supernatants produced by this retroviral vector could, depending on your cloned insert, contain potentially hazardous recombinant virus. Due caution must be exercised in the production and handling of recombinant retrovirus. Appropriate NIH, regional, and institutional guidelines apply.

References

1. Haas, J. *et al.* (1996) *Curr. Biol.* **6**(3):315–324.
2. Gorman, C. (1985). In DNA cloning: A practical approach, vol. II. Ed. D.M. Glover. (IRL Press, Oxford, U.K.) pp. 143–190.
3. Kinsella, T. M. & Nolan G. P. (1996) *Hum. Gene Ther.* **7**(12):1405–1413.
4. Ory, D. S. *et al.* (1996) *Proc. Nat. Acad. Sci. USA* **93**(21):11400–11406.
5. Pear, W. S. *et al.* (1993) *Proc. Natl. Acad. Sci. USA* **90**(18):8392–8396.
6. Yang, S. *et al.* (1999) *Hum. Gene Ther.* **10**(1):123–132.
7. Emerman, M. & Temin, H. M. (1984) *Cell* **39**(3):449–467.

Notice to Purchaser

Clontech products are to be used for research purposes only. They may not be used for any other purpose, including, but not limited to, use in drugs, *in vitro* diagnostic purposes, therapeutics, or in humans. Clontech products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products or to provide a service to third parties without written approval of Clontech Laboratories, Inc.

DH5 α ™ is a trademark of Invitrogen Corporation

Living Colors Products AcGFP1, AmCyan, AsRed, mBanana, mCherry, DsRed, HcRed, mOrange, mPlum, mRaspberry, mStrawberry, tdTomato, ZsGreen, ZsYellow and their variants:

Not-For-Profit Entities: Orders may be placed in the normal manner by contacting your local representative or Clontech Customer Service at 650.919.7300. At its discretion, Clontech grants Not-For-Profit Entities a non-exclusive, personal, limited license to use this product for non-commercial life science research use only. Such license specifically excludes the right to sell or otherwise transfer this product, its components or derivatives thereof to third parties. No modifications to the protein coding sequence may be made without express written permission from Clontech. Any other use of this product requires a license from Clontech. For license information, please contact a licensing representative by phone at 650.919.7320 or by e-mail at licensing@clontech.com.

For-Profit Entities wishing to use this product are required to obtain a license from Clontech. For license information, please contact a licensing representative by phone at 650.919.7320 or by e-mail at licensing@clontech.com or [click here](#) for more information.

This product is the subject of U.S. patents.

For Research Use Only, Not for Use in Humans.

Clontech, the Clontech logo and all other trademarks are the property of Clontech Laboratories, Inc., unless noted otherwise. Clontech is a Takara Bio Company. ©2009 Clontech Laboratories, Inc.